

## Stat Support Activity: Two-Population Mean Confidence Interval Calculations

### Notes: Two-Population Mean Confidence Interval Formula (Independent Groups)

$$(\bar{x}_1 - \bar{x}_2) \pm \left( T_c \times \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)} \right)$$

1.

A random sample of 20 male German Shepherds found that their average weight was 112 pounds with a standard deviation of 28 pounds. A random sample of 14 male Dobermans found that their average weight is 107 pounds with a standard deviation of 24 pounds. Assume that weights are normally distributed. Population 1 is the weight of German Shepherds ( $\mu_1$ ) and population 2 is the weight of Doberman Pincers ( $\mu_2$ ). Use the following formulas to create the following 90% two-population mean confidence interval for the difference between the independent groups ( $\mu_1 - \mu_2$ ).

Sample 1:  $n_1 = 20$  ,  $\bar{x}_1 = 112$  ,  $s_1 = 28$

Sample 2:  $n_2 = 14$  ,  $\bar{x}_2 = 107$  ,  $s_2 = 24$

- Calculate the sample mean difference  $\bar{x}_1 - \bar{x}_2$ .
- Calculate the standard error using the following formula:

$$\text{Standard Error} = \sqrt{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)} =$$

- Calculate the Margin of Error using the following formula and the answers to part (b). (The critical value T-score ( $T_c$ ) = 1.697)

$$\text{Margin of Error} = T_c \times \text{Standard Error}$$

- Calculate the confidence Interval lower limit. Use the answers to part (a) and part (c).

$$(\bar{x}_1 - \bar{x}_2) - \text{Margin of Error}$$

- Calculate the confidence Interval upper limit. Use the answers to part (a) and part (c).

$$(\bar{x}_1 - \bar{x}_2) + \text{Margin of Error}$$

## Stat Support Activity: Two-Population Mean Confidence Interval Calculations

2.

Cotinine is an alkaloid found in tobacco and is used as a biomarker for exposure to cigarette smoke. It is especially useful in examining a person's exposure to second hand smoke. A random sample of 90 non-smoking American adults was collected. These adults were not smokers and did not live with any smokers. The sample mean average cotinine level for this sample was 7.2 ng/mL with a sample standard deviation of 5.8 ng/mL. A second sample of 85 non-smoking American adults was then collected. These adults did not smoke themselves, but did live with one or more smokers. The sample mean average cotinine level for this sample was 28.5 and had a sample standard deviation of 11.4. Population 1 was people that do NOT live with smokers ( $\mu_1$ ) and population 2 was people that DO live with smokers ( $\mu_2$ ). Use the following formulas to create the following 95% two-population mean confidence interval for the difference between the independent groups ( $\mu_1 - \mu_2$ ).

Sample 1:  $n_1 = 90$  ,  $\bar{x}_1 = 7.2$  ,  $s_1 = 5.8$

Sample 2:  $n_2 = 85$  ,  $\bar{x}_2 = 28.5$  ,  $s_2 = 11.4$

- Calculate the sample mean difference  $\bar{x}_1 - \bar{x}_2$ .
- Calculate the standard error using the following formula:

$$\text{Standard Error} = \sqrt{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)} =$$

- Calculate the Margin of Error using the following formula and the answers to part (b). (The critical value T-score ( $T_c$ ) = 1.979)

$$\text{Margin of Error} = T_c \times \text{Standard Error}$$

- Calculate the confidence Interval lower limit. Use the answers to part (a) and part (c).

$$(\bar{x}_1 - \bar{x}_2) - \text{Margin of Error}$$

- Calculate the confidence Interval upper limit. Use the answers to part (a) and part (c).

$$(\bar{x}_1 - \bar{x}_2) + \text{Margin of Error}$$